



T R A C I E

HEALTHCARE EMERGENCY PREPAREDNESS
INFORMATION GATEWAY

Access the webinar here:

<https://attendee.gotowebinar.com/recording/9109707139001760272>

Access speaker bios here:

<https://files.asprtracie.hhs.gov/documents/healthcare-operations-speaker-series-bios--respiratory-therapists.pdf>

Access the entire webinar series here:

<https://files.asprtracie.hhs.gov/documents/lessons-learned-in-healthcare-operations-during-a-pandemic-speaker-series-summary.pdf>

Healthcare Operations during the COVID-19 Pandemic- Speaker Series

May 2021

Unclassified//For Public Use



The Role of the AARC in Pandemic Response

American Association for Respiratory Care
Irving, TX

Presenter



Douglas S. Laher MBA, CAE, RRT, FAARC
Chief Operating Officer
American Association for Respiratory Care

Objectives

- Who is the respiratory therapist (RT)?
- Education, credentialing, & licensure
- RT specialties
- Who is the AARC?
- Response of the AARC during the pandemic

Allied health professional that specialize in managing patients having ventilation and/or oxygenation disorders

- Clinical focus:
 - Diagnostic evaluation
 - Initial and subsequent evaluation of patient condition and response to plan of care
 - Patient education
 - Disease management
 - Disease prevention



Education, Credentialing & Licensure

Training

- Associate degree
- Bachelor's degree
- Master's degree



Credentialed

- National Board for Respiratory Care
 - Certified Respiratory Therapist (CRT)
 - Registered Respiratory Therapists (RRT)
 - Certified Pulmonary Function Technologist (CPFT)
 - Registered Pulmonary Function Technologist (RPFT)
 - Specialty credentials in adult acute care, neonatal/pediatrics, and sleep



Licensed in 49 states



RT Specialties

Adult Acute
Care

Ambulatory &
Post-Acute
Care

Education

Leadership &
Management

Neonatal-
Pediatrics

Diagnostics

Sleep

Surface & Air
Transport



American Association
for Respiratory Care

Who is the AARC?

- Only national 501-c6 Association representing the Respiratory Therapist
- More than 40,000 members across the globe
- Membership from 44 different countries
 - 50 state affiliates in U.S.
 - 6 international affiliates
- AARC Mission Statement

“The AARC is the foremost professional association promoting respiratory therapists.”
- AARC Vision Statement

“The AARC advances professional excellence and science in the practice of respiratory therapy, serving the profession, patients, caregivers and the public.”

AARC's Response to Pandemic

- Communication
- Guidance Documents/Joint Statements
- Peer-reviewed Publications Education
- Media Outreach
- Advocacy
- COVID-19 RT Heroes Fund



A screenshot of the AARC website. The top navigation bar includes the AARC logo, contact information (9425 North Heather Blvd., Suite 100, Irving, TX 75063), and a CARE number (75063-4706). The main content area features a large blue heart graphic with yellow wings, and the text 'COVID-19 RT Fund'. Below this, there is a section titled 'U.S. States and Territories Modifying Requirements for Telehealth in Response to COVID-19' with a table of states and territories. The table lists states with waivers, states with waivers not allowing new regulations, and states without waivers. The bottom section of the screenshot shows a 'Respiratory therapists are keeping ventilators alive' article, with a sub-header 'Doctors know these professionals are essential'.

The Role of the Respiratory Therapist in Pandemic Response

Rush University Medical Center
Rush University
Chicago, Illinois



Rush Respiratory Care



J. Brady Scott, PhD, RRT, RRT-ACCS, AE-C, FAARC, FCCP

Director of Clinical Education, Associate Professor

Department of Cardiopulmonary Sciences, Division of Respiratory Care, College of Health Sciences, Rush University



David L. Vines, PhD, MHS, RRT, FAARC, FCCP


Chairperson, Academic Program Director, Associate Professor

Department of Cardiopulmonary Sciences, Division of Respiratory Care, College of Health Sciences, Rush University

Objectives

- Describe the training and skillset of the Respiratory Therapist to be leveraged in disaster response
- Apply examples from the COVID-19 pandemic to the role of the Respiratory Therapist in disaster response

Respiratory Therapy in Pandemic Response – Pre-Pandemic Surge



Rapid Training



Evaluation of a Training Method to Improve Knowledge and Confidence of Prone Positioning

Afrah Obaidan, MSc, RRT, RRT-NPS

J. Brady Scott, MSc, RRT, RRT- ACCS, AE-C, FAARC

Sara Hanif Mirza, MD, MS

Adel Aljoaid, MSc, RRT, RRT-ACCS

Rima Tailor, MSc, RRT, RRT-NPS

David L. Vines, MHS, RRT, FAARC

Conclusion

In institutions where these resources are available discussion, video demonstration, and simulation with standardized patients may increase ICU clinician knowledge and confidence with prone positioning.



Appendix C

Protocol for prone positioning

Objective

- To establish safety measures for prone positioning of mechanically ventilated patients

Policy

- Prone positioning may be used in the intensive care unit in an attempt to improve arterial oxygenation and pulmonary mechanics in patients with acute lung injury or acute respiratory distress syndrome (ARDS).
- Prone may be performed manually or with the Roto-Prone™ Therapy System bed, which may be ordered by an ICU physician.

Indications

Patients who meet the following criteria:

- Endotracheal intubation and mechanical ventilation for ARDS for less than 36 hours
- Severe ARDS defined as :
 - PaO₂:FIO₂ ratio of <150 mm Hg
 - FIO₂ of ≥ 0.6
 - PEEP of ≥10 cm H₂O of water

Contraindications

- Increased intracranial pressure (ICP) >30 mm Hg or cerebral perfusion pressure <60 mmHg (CPP = MAP-ICP)
- Recent tracheal surgery or sternotomy
- Recent facial trauma or surgery
- Spine instability
- Deep venous thrombosis treated for less than 2 days
- Recent cardiac pacemaker insertion
- Unstable bone fractures
- Hemodynamic instability; mean arterial pressure <65 mm Hg
- Pregnancy
- Chest tube with air leaks (anterior)
- Patient weight > 350 lbs

Procedure

REQUIRES THE FOLLOWING STAFF:

a. Registered nurses (prepare the patient)

- Securing the lines, tubes, and drains
- Monitoring vital signs
- Turning of the patient
- Positioning of the patient

NOTE: An inflatable mattress or air bed is preferred, but not required, for the procedure.

b. Respiratory therapist

- Suctioning
- Ventilator manipulation
- General assistance

c. Physician (Available if problems arise)

NOTE: Other staff may be needed depending on the size of the patient.

Guidelines for the prone positioning:

- People required: 3-4 (one person dedicated to the management of the head, endotracheal tube, and ventilator circuit).
- The person at the head of the bed coordinates the steps of the procedure and secures the airway.
- The other people stand on either side of the bed.
- The decision for which direction for the rotation should give priority to the side of the central lines (central lines go upward rather than rolled on to).
- Check that the length of the vascular lines and the ventilator circuit is appropriate in order to prevent tension during the turn.
- Endotracheal and gastric tubes must be secured.
- The patient is moved horizontally to the side opposite the direction of the rotation.
- The patient is moved in the sagittal plane and kept there while the cardiac electrodes are moved to their back and a new bed sheet is set.
- The patient is rotated to the full prone position and the upper limbs are placed alongside the body.
- Place pillows for positioning under the patient's chest, pelvis, and lower legs.

Appendix D Prone positioning checklist

Turning	
<input type="checkbox"/>	People required: 3-4 (one person dedicated to the management of the head, endotracheal tube, and ventilator circuit).
<input type="checkbox"/>	The person at the head of the bed coordinates the steps of the procedure.
<input type="checkbox"/>	The other people stand on either side of the bed.
<input type="checkbox"/>	The decision for which direction for the rotation gives priority to the side of the central lines (central lines go upward rather than rolled on to).
<input type="checkbox"/>	Check that the vascular lines and ventilator circuit length are appropriate.
<input type="checkbox"/>	Endotracheal and gastric tubes are secured.
<input type="checkbox"/>	The patient is moved horizontally to the side opposite the direction of the rotation.
<input type="checkbox"/>	The patient is moved in the sagittal plane and kept there while the cardiac electrodes are moved to their back and a new bed sheet is set.
<input type="checkbox"/>	The patient is rotated to the full prone position and the upper limbs are placed alongside the body.
<input type="checkbox"/>	Place pillows for positioning under the patient's chest, pelvis, and lower legs.
Immediately after the turn	
<input type="checkbox"/>	Turn the head and neck of the patient alternately from left to right every 2 hours.
<input type="checkbox"/>	Reverse Trendelenberg: slight degrees of reverse Trendelenberg (10 - 20 degrees) are often well tolerated and may be useful in certain patients during prone positioning.
<input type="checkbox"/>	Leave the patient in prone position up to 16 hours, then 2-4 hours in the supine position.
<input type="checkbox"/>	Reassess the security and patency of all tubes/lines.
<input type="checkbox"/>	ETT distance Cuff leak Pressure points around ETT and securement device Check for any kinks in tubing Breath sounds, ventilator parameters Lifting team to assist RRT to establish airway patency. The head and shoulders may need to be lifted and supported in order for ventilator tubing to hang freely.
<input type="checkbox"/>	Reassess SpO ₂ , blood pressure, cardiac rhythm, and breath sounds.
<input type="checkbox"/>	Reassess ETT/Trach leak. (May adjust cuff volume, head position, and delivered Vt to assure adequate ventilation.)
<input type="checkbox"/>	Uncap/reattach capped off lines/NGT/NJT.

Prone Positioning for ARDS



Prone Positioning for Acute Respiratory Distress Syndrome (ARDS)

452,414 views · Apr 2, 2020

2.3K

129


SHARE

SAVE

...

YouTube: <https://youtu.be/lcBPQHQUvXY>

Respiratory Therapy in Pandemic Response – Pandemic Surge



Interdisciplinary Prone Team

- Respiratory Therapists
- Nurses
- Physical & Occupational Therapists



Respiratory Care Students and COVID-19



Kimberly Villanueva, left, takes part in intubating a mannequin with other students during a class at Rush University Medical Center on July 31, 2020. (Antonio Perez / Chicago Tribune)

High Acuity Interventions for COVID-19

Mechanical Ventilation

- Focus on adequate oxygenation, ventilation, & lung protection
- Protocol utilization
- Monitor and prevent ETT occlusions


Prone Positioning

- Improve oxygenation & promote lung protective mechanical ventilation
- Awake patients on noninvasive support

Noninvasive Support (HFNC/NIV)

- Reduce the need for invasive mechanical ventilation
- Allows for care outside of full ICUs

Respiratory Therapy in Pandemic Response – Post-Pandemic Surge



Prone positioning for patients intubated for severe acute respiratory distress syndrome (ARDS) secondary to COVID-19: a retrospective observational cohort study

Tyler T. Weiss¹, Flor Cerda², J. Brady Scott^{1,3}, Ramandeep Kaur¹, Sarah Sungurlu⁴, Sara H. Mirza^{1,4}, Amnah A. Alolaiwat³, Ramandeep Kaur³, Ashley E. Augustynovich³ and Jie Li^{1,3,*}

¹Department of Respiratory Care, Rush University Medical Center, Chicago, IL, USA, ²Department of Nursing, Medical Intensive Care Unit, Rush University Medical Center, Chicago, IL, USA, ³Department of Cardiopulmonary Sciences, Division of Respiratory Care, Rush University, Chicago, IL, USA and ⁴Department of Pulmonary and Critical Care, Rush University Medical Center, Chicago, IL, USA

*Corresponding author. E-mail: jie_li@rush.edu

In summary, prone positioning improved oxygenation for patients with COVID-19 ARDS who required invasive mechanical ventilation. Serial assessment of the P_{aO_2}/F_{iO_2} ratio may help guide decisions for earlier escalation of treatment, including ECMO.

Original Clinical Report

Critical Care Explorations

Effects of Inhaled Epoprostenol and Prone Positioning in Intubated Coronavirus Disease 2019 Patients With Refractory Hypoxemia

Jie Li, PhD, RRT, RRT-ACCS, RRT-NPS¹; James B. Fink, PhD, RRT^{1,2}; Ashley E. Augustynovich, MS, RRT¹; Sara Mirza, MD, MS^{1,3}; Richard H. Kallet, MS, RRT⁴; Rajiv Dhand, MD⁵

CONCLUSIONS

In mechanically ventilated patients with COVID-19 who had refractory hypoxemia, combined use of iEPO and PP improved oxygenation to a greater extent than with each treatment individually. Responders to combined modalities had lower mortality than nonresponders.

Airborne Particulate Concentrations During and After Pulmonary Function Testing

To the Editor:

Pulmonary function tests (PFTs) are an integral component of the evaluation of patients with pulmonary diseases.¹ Due to concerns for virus transmission, multiple respiratory societies



recommend to postpone or limit PFTs during the coronavirus disease 2019 pandemic.²⁻⁶ Repeated forced breathing maneuvers during PFTs may generate bioaerosol by airway opening^{7,8} or inducing cough.^{1,7} However, the concentrations of particles that are generated and change over time during and after PFTs are unknown, leading to the current recommendation to close PFT laboratories for 20 minutes to 3 hours between tests.²⁻⁴ Thus, we investigated aerosol particle generation and clearance during and after PFTs.

[159 #4 CHEST APRIL 2021]

Although performing PFTs in negative-pressure rooms may be preferred, our data suggest that reductions of ambient particles can be achieved in rooms with less aggressive ventilation exchanges and that exposure to staff members during and after PFT procedures is, to some extent, independent of the particle clearance time. To avoid transmission of infection, PFT technologists should take high-level personal protective equipment precautions during testing of any patient during this pandemic. Alternative methods that include portable electronic spirometry and self-monitoring technologies might be considered.¹¹

Kaur et al. *Critical Care* (2020) 24:571
<https://doi.org/10.1186/s13054-020-03231-8>


Critical Care

REVIEW

Open Access



Practical strategies to reduce nosocomial transmission to healthcare professionals providing respiratory care to patients with COVID-19

Ramandeep Kaur¹, Tyler T. Weiss¹, Andrew Perez¹, James B. Fink¹, Rongchang Chen², Fengming Luo³, Zongan Liang³, Sara Mirza¹ and Jie Li^{1*} 

Conclusion

The frontline HCWs are at risk for contracting the COVID-19 disease when caring for patients and providing aerosol-generating procedures. Until further high-quality studies generate robust evidence, defining the precise nosocomial transmission risk associated with AGMPs, along with CDC's recommended PPE guidelines, we propose additional respiratory protective measures that could reduce the nosocomial transmission of COVID-19 diseases to HCWs providing respiratory interventions.

PULMONARY PERSPECTIVE

Coughs and Sneezes: Their Role in Transmission of Respiratory Viral Infections, Including SARS-CoV-2

✉ Rajiv Dhand¹ and Jie Li²

¹Division of Pulmonary and Critical Care Medicine, Department of Medicine, University of Tennessee Graduate School of Medicine, Knoxville, Tennessee; and ²Division of Respiratory Care, Department of Cardiopulmonary Sciences, Rush University Medical Center, Chicago, Illinois

ORCID IDs: 0000-0002-3621-2422 (R.D.); 0000-0003-0121-1291 (J.L.).

American Journal of Respiratory and Critical Care Medicine Volume 202 Number 5 | September 1 2020



Conclusions

Coughs and sneezes create respiratory droplets of variable size that spread respiratory viral infections. Because these droplets are forcefully expelled, they are dispersed in the environment and can be inhaled by a susceptible host. Whereas most respiratory droplets are filtered by the nose or deposit in the oropharynx, the smaller droplet nuclei become suspended in room air and individuals farther away from the patient could inhale them. These finer

Open access

Protocol

BMJ Open Awake prone positioning of hypoxaemic patients with COVID-19: protocol for a randomised controlled open-label superiority meta-trial

Elsa Tavernier ^{1,2} Bairbre McNicholas,^{3,4} Ivan Pavlov,⁵ Oriol Roca,^{6,7} Yonatan Perez,⁸ John Laffey,^{3,4} Sara Mirza,⁹ David Cosgrave,^{3,4} David Vines,⁹ Jean-Pierre Frat,¹⁰ Stephan Ehrmann,⁸ Jie Li ⁹

Tavernier E, et al. *BMJ Open* 2020;**10**:e041520. doi:10.1136/bmjopen-2020-041520

Strengths and limitations of this study

- ▶ This pragmatic design will deal with the recruitment difficulties that could occur in the individual trials given the uncertainties of the international dynamics of the COVID-19 pandemic.
- ▶ The collaborative interim analysis plan at the level of the meta-trial will enable an earlier data analysis compared with the individual study level or to a retrospective meta-analysis.
- ▶ Besides synthesising the effect size estimates, it also considers the aspect of replication: results being consistent across trials is a strength in favour of a robust treatment effect over different conditions.
- ▶ The lack of blinding of trial participants, care providers and outcome assessors is an unavoidable limitation of the study design.

USNews NEWS » News Best Countries Best States Healthiest Communities Cities Elections The Racial Divide

Home / News / Healthiest Communities

COMMENTARY

Even After the Coronavirus Pandemic, America Can't Breathe Easy

It's crucial that the U.S. ramp up its workforce of respiratory therapists now, but a need for more providers will still exist in the wake of COVID-19.

By J. Brady Scott | April 1, 2020, at 10:00 a.m.

The Washington Post
Democracy Dies in Darkness

Sections

Coronavirus U.S. map World map Vaccine tracker Vaccine FAQ Variants FAQ A pandemic year Coronavirus Living

Home

Share

7

PostEverything • Perspective

Respiratory therapists are keeping many covid patients on ventilators alive

Doctors know these professionals are essential



Low Tech Way to Help Some Covid Patients: Flip Them Over *New York Times*
<https://www.nytimes.com/2020/05/13/health/coronavirus-proning-lungs.html>

Media



CORONAVIRUS PANDEMIC

U.S.		WORLD	
CONFIRMED	DEATHS	CONFIRMED	DEATHS
671,425	33,286	2,172,031	146,291

SOURCE: WHO, CDC, ECDC, NHC, DXY

U.S. STATES	CASES	DEATHS
ILLINOIS	25,733	1,072
FLORIDA	23,340	668
LOUISIANA	22,532	1,156

DOW WATCH

24,044.30
▲ 506.62 2.15%


FOX NEWS channel

HEALTH CARE WORKERS TREATING COVID-19 PATIENTS SHARE THEIR STORIES WITH FOX NEWS

AMERICA'S NEWSROOM

SEN KELLY LOEFFLER (R-GA)
10:30 AM ET

This Morning



CORONAVIRUS PANDEMIC

IN THE UNITED STATES

TOTAL CASES
15,618,685

DEATHS
292,192

SOURCE: JOHNS HOPKINS UNIVERSITY

SUNDAY ON CNN
CNN HEROES
HOSTED BY ANDERSON COOPER & KELLY RIPA
8 P. ET

CORONAVIRUS PANDEMIC

FRONTLINE WORKERS TO RECEIVE VACCINE FIRST AS HOSPITALIZATIONS SURGE

Brady Scott | Respiratory Therapist, Rush University Medical Center

LIVE CNN

DOW ▲ 125.66

Respiratory Therapists and Faculty at RUMC



Special Acknowledgements:

Department of Cardiopulmonary Sciences, Division of Respiratory Care
Respiratory Care Services, RUMC

Steve Mosakowski, MBA, RRT, RRT ACCS,
RRT NPS, CPFT, FAARC Director of Clinical
Services, Assistant Professor

Ellen Becker, PhD, RRT, RRT NPS, RPFT, AE
C, FAARC, Professor

Jie Li, PhD, RRT, RRT ACCS,
RRT NPS, FAARC, Associate Professor

Ramandeep Kaur, PhD, RRT, RRT ACCS, AC
E, Respiratory Therapist 3 / Research coordinator,
Assistant Professor

Tyler Weiss, MSc, RRT, RRT ACCS, AC E
Education Coordinator, Assistant Professor

Ankeet Patel, MSc, RRT, RRT ACCS Clinical
Manager, Adult Care Section, Assistant professor

Andrew Klein, MSc, RRT, RRT ACCS, RRT
NPS, AC E, Respiratory Therapist 3, Adult Clinical
Specialist

Ahmad Elshafei, MSc, RRT, RRT ACCS, RRT
NPS, AC E, Respiratory Therapist 3, Adult Clinical
Specialist

Ashley Lachowicz, MSc, RRT, RRT-ACCS,
RRT-NPS, AC-E, Respiratory Therapist 3,
Adult Clinical Specialist

Lauren Harnois, MSc, RRT, RRT-ACCS,,
Respiratory Therapist 3, Adult Clinical
Specialist

Anne Geistkemper, MSc, RRT, RRT
NPS, Clinical Manager, Neonatal Pediatric
Section, Instructor

Sara Murphy, MBA, RRT, RRT NPS.
RRT ACCS, AE C, Clinical Educator for
Neonatal Pediatric Section, Instructor

Jacob Burd, MSc, RRT, RRT NPS., C
NPT, Respiratory Therapist 3, Neonatal
Pediatric Specialist

Katie Dugan, MSc, RRT, RRT NPS.,
Respiratory Therapist 3, Neonatal Pediatric
Specialist

The entire Respiratory Care staff at Rush University Medical Center

Thank you.

Rush is a not-for-profit health care, education and research enterprise established in Chicago, Illinois in 1837.



Colleges of Medicine, Nursing, Health Sciences and Graduate College



Contact ASPR TRACIE



asprtracie.hhs.gov



1-844-5-TRACIE



askasprtracie@hhs.gov